

Single-electron transport by surface acoustic waves

Vyacheslavs Kashcheyevs

*School of Physics and Astronomy,
Raymond and Beverly Sackler Faculty of Exact Sciences
Tel Aviv University, Tel Aviv 69978, ISRAEL*

We use the theory of single electron adiabatic quantum pumping to calculate the current driven through a depleted quantum point contact by a surface acoustic wave (SAW). For sufficient SAW power, an (almost) integer number of electrons is transferred per SAW period, thus giving quantized values to the current [1, 2]. This quantization is a result of quantum interference and does not require a Coulomb-blockade effect. We reproduce qualitatively various experimentally observed properties of the quantized acoustoelectric current [3], which include the dependance on SAW amplitude and wavelength, accuracy enhancement by a counter-propagating SAW and the effects of an additional source-drain bias. Conditions for optimal single-electron transport are discussed.

-
- [1] A. Aharony and O. Entin-Wohlman, Phys. Rev. B **65**, 241401 (2002), con-dmat/0111053.
 - [2] O. Entin-Wohlman, A. Aharony, and V. Kashcheyevs, J. Phys. Soc. Japan **14A**, 77 (2003), cond-mat/0201073.
 - [3] J. Cunningham, V. I. Talyanskii, J. M. Shilton, M. Pepper, M. Y. Simmons, and D. A. Ritchie, Phys. Rev. B **60**, 4850 (1999), and references there-in.